

Specimen Label



Gallery*

75 Dry Flowable

Specialty Herbicide

*Trademark of Dow AgroSciences LLC

A selective preemergence herbicide for control of certain broadleaf weeds in:

- Established turf
- Landscape ornamentals
- Container grown ornamentals
- Nursery stock
- Ground covers
- Ornamental bulbs
- Non-bearing fruit and nut trees
- Christmas tree plantations and non-bearing vineyards
- Non-cropland

Active Ingredient:

isoxaben: *N*-[3-(1-ethyl-1-methylpropyl)-5-isoxazoly]-2,6-dimethoxybenzamide and isomers 75%

Inert Ingredients 25%
Total 100%

Contains 0.75 pounds active ingredient per pound.

EPA Reg. No. 62719-145

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes Eye Irritation • Harmful If Inhaled

Avoid ingestion, breathing dust or spray mist, and contact with skin, eyes, or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes Flush eyes with plenty of water. Call a physician if irritation develops.

If swallowed Call a physician or Poison Control Center.

If on skin Wash with plenty of soap and water. Get medical attention if irritation develops.

If inhaled: Remove individual to fresh air. If not breathing, provide cardiopulmonary resuscitation assistance and get medical attention.

Environmental Hazards

Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift may result in reduced germination or emergence of non-target plants adjacent to treated area.

Notice: Read the entire label. Use only according to label directions.

Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" elsewhere on this label.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.


Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

NOTICE: The container cap for Gallery* 75 Dry Flowable herbicide may be used to measure the amount of product required to treat small areas. When filled to the level of the inner lip, the cap contains approximately 1/2 ounce which is the amount of Gallery 75 Dry Flowable needed to treat an area of 1,000 square feet. Please refer to label booklet for complete use directions and precautions.



When filled to this level, container cap contains (14 grams) of Gallery 75 Dry Flowable.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements†

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exemptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container. Do not store in direct sunlight. Do not store at temperatures above 120°F. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Dispose of in a sanitary landfill, or incineration, or by burning if allowed by state and local authorities. If burned, stay out of smoke.

General Information

Gallery 75 Dry Flowable is a preemergence herbicide for control of certain broadleaf weeds in established turf, landscape ornamentals, container grown ornamentals, nursery stock, ground covers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards, Christmas tree plantations and non-cropland areas. Apply Gallery 75 Dry Flowable in late summer to early fall, in early spring, or anytime prior to germination of target weeds, or immediately after cultivation.

Gallery 75 Dry Flowable does not control established weeds. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings, and trash should be removed or thoroughly mixed into the soil prior to treatment. Soil in non-turf areas should be in good condition and free of clods at the time of application. Gallery 75 Dry Flowable is stable on the soil surface for up to 21 days, but must be activated to be effective. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation after application is necessary to activate Gallery 75 Dry Flowable. If Gallery 75 Dry Flowable is not activated by rainfall or irrigation within 21 days of application, erratic weed control may result. In non-turf areas, if weeds emerge due to lack or rainfall or irrigation, shallow cultivate to a depth of 1 to 2 inches to destroy existing weeds or remove by hand.

In Arizona: The state of Arizona has not approved Gallery 75 Dry Flowable for use on plants grown for commercial production such as: turf sod farms, ornamentals and nursery stock grown for resale, and non-bearing fruit and nut trees and non-bearing vineyards.

Mixing and Application Directions

Gallery 75 Dry Flowable - Alone

Check to be sure spray equipment is clean and not contaminated. Using clean water, fill the tank to 1/2 of the final volume required and start agitation. Slowly add the correct quantity of Gallery 75 Dry Flowable to the spray tank, continue agitation and complete filling the tank. Maintain agitation during filling and through application. Agitation should be sufficient to create a rippling or rolling action on the liquid surface. Sparger pipe agitation generally provides the best agitation. In-line screens and strainers should be no finer than 16 mesh.

If spraying and agitation is stopped, Gallery 75 Dry Flowable may settle to the bottom of the spray tank. If settling occurs, material must be re-suspended before continuing spray application. Clean the spray tank, lines and screens thoroughly after use.

Gallery 75 Dry Flowable in Tank Mixes

Add Gallery 75 Dry Flowable to the spray tank as described above and fill. Fill the spray tank to 3/4 of the final volume required. Add other formulations in this order: (1) other dry flowables, (2) wettable powders, (3) aqueous suspensions, (4) flowables, (5) liquids, (6) solutions and emulsifiable or liquid concentrates. Allow two to three minutes between the addition of each product, and agitate continuously until the product(s) are completely dispersed in water. Maintain agitation during filling and through application. If a build-up of materials is observed on the walls of the spray tank, wash the tank with soapy water between fillings, rinse and then continue the spraying operation. Follow label directions for each material added to the tank.

Application

Apply Gallery 75 Dry Flowable at 0.66 to 1.33 lb/acre with a properly calibrated low pressure herbicide sprayer that provides uniform distribution. Nozzle screens should be no finer than 50 mesh (50 mesh is finer than 16 mesh). In-line screens and strainers should be no finer than 16 mesh. Apply Gallery 75 Dry Flowable in 10 to 200 gallons of water carrier per acre. As the spray volume decreases, the importance of accurate calibration and uniform application increases. Take precautions to avoid spray drift when applying Gallery 75 Dry Flowable. Drift may result in reduced germination or emergence of non-target plants adjacent to the treated area. Maintain agitation from mixing through application. Avoid boom overlaps that will increase rates above those recommended. Calibrate application equipment prior to use, according to manufacturer's directions. Check calibration frequently to be sure equipment is working properly and distributing spray uniformly.

NOTICE: The container cap for Gallery 75 Dry Flowable herbicide may be used to measure the amount of product required to treat small areas. When filled to the level of the inner lip, the cap contains approximately 1/2 ounce which is the amount of Gallery 75 Dry Flowable needed to treat an area of 1,000 square feet. (See illustration at front of booklet.)

Weeds Controlled or Suppressed by Gallery 75 Dry Flowable

Weeds controlled when applied at 0.66 lb/acre (0.25 oz or 7 grams/1000 sq ft)

Common Name

aster, slender
bursage, annual
celery, wild
chickweed, common
clover, white
cudweed, purple
fiddleneck, coast
filaree, redstem
fleabane, blackleaved
fleabane, dwarf

Scientific Name

Aster exilis
Ambrosia acanthicarpa
Apium leptophyllum
Stellaria media
Trifolium repens
Gnaphalium purpureum
Amsinckia intermedia
Erodium cicutarium
Conyza bonariensis
Conyza ramosissima

groundcherry, lanceleaf
 henbit
 horseweed
 knotweed, prostrate
 lambsquarters, common
 mallow, little
 mustard, Indian
 mustard, wild
 nightshade, black
 pepperweed, Virginia
 pigweed
 pineappleweed
 plantain, slender
 purslane, common
 radish, wild
 ragweed, common
 rocket, London
 rockpurslane, desert
 shepherdspurse
 sibara
 smartweed, Pennsylvania
 sowthistle, annual
 speedwell, purslane
 spurge, spotted
 telegraphplant
 thistle, Russian
 velvetleaf

Physalis lanceifolia
Lamium amplexicaule
Conyza canadensis
Polygonum aviculare
Chenopodium album
Malva parviflora
Brassica juncea
Sinapis arvensis
Solanum nigrum
Lepidium virginicum
Amaranthus spp.
Matricaria matricarioides
Plantago elongata
Portulaca oleracea
Raphanus raphanistrum
Ambrosia artemisiifolia
Sisymbrium irio
Calandrinia ciliata
Capsella bursa-pastoris
Sibara virginica
Polygonum pensylvanicum
Sonchus oleraceus
Veronica peregrina
Euphorbia maculata
Heterotheca grandiflora
Salsola iberica
Abutilon theophrasti

In addition to the weeds listed at the 0.66 lb/acre and 1.0 lb/acre rates, the following weeds will be controlled at 1.33 lb/acre (0.5 oz or 14 grams/1000 sq ft)

Common Name
 burclover, California
 datura
 dogfennel
 eveningprimrose
 fescue, rattail
 filaree, whitestem
 goosefoot, nettleleaf
 jimsonweed
 knotweed, silversheath
 kochia
 medic, black
 mullein, turkey
 nettle, burning
 oxtongue, bristly
 pimpernel, scarlet
 sowthistle, spiny
 spurge, petty
 spurge, prostrate
 sunflower
 swinecress
 thistle, musk
 willoweed, panicle
 woodsorrel, creeping

Scientific Name
Medicago polymorpha
Datura spp.
Eupatorium capillifolium
Oenothera spp.
Vulpia myuros
Erodium moschatum
Chenopodium murale
Datura stramonium
Polygonum argyrocoleon
Kochia scoparia
Medicago lupulina
Eremocarpus setigerus
Urtica urens
Picris echioides
Anagallis arvensis
Sonchus asper
Euphorbia peplus
Euphorbia humistrata
Helianthus spp.
Coronopus didymus
Carduus nutans
Epilobium paniculatum
Oxalis corniculata

In addition to the weeds listed at the 0.66 lb/acre rate, the following weeds will be controlled at 1.0 lb/acre (0.38 oz or 10.5 grams/1000 sq ft)

Common Name
 aster, heath
 bittercress
 bittercress, hairy
 brassbuttons, southern
 carrot, wild
 chamberbitter
 chickweed, mouseear
 dandelion
 eclipta
 galinsoga, hairy
 geranium, Carolina
 groundsel, common
 ladythumb
 lettuce, prickly
 mallow, dwarf
 marestail
 mayweed
 morningglory, ivyleaf
 mustard, black
 pennywort
 phyllanthus, long-stalk
 plantain, bracted
 plantain, broadleaf
 plantain, buckhorn
 pokeweed, common
 rockpurslane, redmaids

 sida, prickly
 sorrell, red
 speedwell, thymeleaf
 spurge, hyssop
 sweetclover, yellow
 tansymustard, green

 woodsorrel, yellow

Scientific Name
Aster ericoides
Cardamine oligosperma
Cardamine hirsuta
Cotula australis
Daucus carota
Phyllanthus urinaria
Cerastium vulgatum
Taraxacum officinale
Eclipta prostrata
Galinsoga ciliata
Geranium carolinianum
Senecio vulgaris
Polygonum persicaria
Lactuca serriola
Malva rotundifolia
Hippuris vulgaris
Anthemis cotula
Ipomoea hederacea
Brassica nigra
Hydrocotyle spp.
Phyllanthus tenellus
Plantago aristata
Plantago major
Plantago lanceolata
Phytolacca americana
Calandrinia ciliata
 var. menziesii
Sida spinosa
Rumex acetosella
Veronica serpyllifolia
Euphorbia hyssopifolia
Melilotus officinalis
Descurainia pinnata
 spp. *brachycarpa*
Oxalis stricta

In addition to the weeds controlled, the following weeds will be partially controlled or suppressed at 1.33 lb/acre (0.5 oz or 14 grams/1000 sq ft)

Common Name
 bindweed, field
 carpetweed
 dock, curly
 mallow, Venice
 milkweed, honeyvine
 morningglory, tall
 pusley, Florida

Scientific Name
Convolvulus arvensis
Mollugo verticillata
Rumex crispus
Hibiscus trionum
Ampelamus albidus
Ipomoea purpurea
Richardia scabra

Approved Uses

Established Turfgrasses

Gallery 75 Dry Flowable is recommended as a preemergence treatment for control of certain broadleaf weeds in established cool season and warm season turfgrasses.

Apply Gallery 75 Dry Flowable in late summer to early fall, in early spring, or anytime prior to germination of target weeds.

Refer to "General Information" section of this label prior to using this product on turf.

Special Use Precautions

Apply Gallery 75 Dry Flowable only on established turf. Apply to newly seeded turf (including overseeded turf) only after seedlings are established (three leaf stage and tillering) and well rooted. Do not overseed established turf sooner than 60 days following an application of Gallery 75 Dry Flowable.

Do not use Gallery 75 Dry Flowable on golf course putting greens or tees.

Do not use Gallery 75 Dry Flowable on dichondra.

Do not use Gallery 75 Dry Flowable on grass grown for seed.

Treatment of Turf Species Not Listed on the Label for Gallery 75 Dry Flowable

Users who wish to use Gallery 75 Dry Flowable on turf species not recommended on this label may determine the suitability for such uses by treating a small area at a recommended rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target turf species. **The user assumes the responsibility for any turf damage or other liability resulting from use of Gallery 75 Dry Flowable on turf species not recommended on this label.**

Tank Mixing

Gallery 75 Dry Flowable may be tank mixed with Surflan* A.S. and applied to **warm season turfgrass only** as a preemergence treatment to provide broad spectrum annual grass and broadleaf weed control. Gallery 75 Dry Flowable may also be applied as a separate treatment to supplement Balan 2.5G or Team 2G applications in cool and warm season turf. Gallery 75 Dry Flowable may be tank mixed with postemergence broadleaf herbicides registered for use on established turf to control existing broadleaf weeds and provide residual preemergence broadleaf weed control. Applied as directed, Gallery 75 Dry Flowable tank mixes will provide control of susceptible weed species listed on the respective labels. When using Gallery 75 Dry Flowable in tank mix combinations with other products, read and follow all applicable use directions, precautions, and limitations on the respective product labels.

Gallery 75 Dry Flowable is recommended for use on the following turf species:

Established Cool Season Turf Grasses

Common Name	Scientific Name
bentgrass, creeping	<i>Agrostis stolonifera</i>
bentgrass, colonial	<i>Agrostis tenuis</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewing	<i>Festuca rubra</i> var. <i>commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheeps	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

Established Warm Season Turf Grasses

Common Name	Scientific Name
bahiagrass	<i>Paspalum notatum</i> var. <i>Saurae parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
buffalograss	<i>Buchloe dactyloides</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

Landscape Ornamentals, Container Grown Ornamentals, Nursery Stock, Ground Covers, Ornamental Bulbs, Non-bearing† Fruit and Nut Trees and Non-bearing† Vineyards, and Christmas Tree Plantations

†Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment. **Gallery 75 Dry Flowable is recommended as a preemergence treatment for control of certain broadleaf weeds in landscape ornamentals, container grown ornamentals, nursery stock, ground covers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards, and Christmas tree plantations.**

Apply Gallery 75 Dry Flowable anytime prior to germination of target weeds, or immediately after cultivation.

Note: Refer to the "General Information" section of this label for use precautions and information on mixing and application, application rates, and weeds controlled prior to using this product.

Special Use Precautions

To avoid possible injury, do not apply Gallery 75 Dry Flowable to: Nursery seedbeds or forest or Christmas tree seedling transplant beds.

Unrooted liners or cuttings that have been planted in pots for the first time.

Pots less than four inches wide.

Bedding plants or areas where bedding plants will be planted or transplanted within one year following application.

Ground covers until they are established and well rooted.

Do not apply Gallery 75 Dry Flowable to newly transplanted ornamentals, nursery stock, ground covers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards or Christmas trees until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or plant injury may occur.

When planting into a site treated with Gallery 75 Dry Flowable in the past 8 months, use untreated soil as fill around roots when replacing plants or injury may occur.

Repeat applications at 1.0 pound per acre or more of Gallery 75 Dry Flowable should not be made sooner than 60 days after a previous application of Gallery 75 Dry Flowable. Do not apply more than a total of 4.0 pounds per acre of Gallery 75 Dry Flowable within a 12 month period.

Treatment of Plant Species Not Listed on the Gallery 75 Dry Flowable Label

Users who wish to use Gallery 75 Dry Flowable on plant species not recommended on this label may determine the suitability for such uses by making a trial application of Gallery 75 Dry Flowable at a recommended rate to a small number of plants. Prior to using Gallery 75 Dry Flowable on a larger number of plants, the treated plants should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. **The user assumes the responsibility for any plant damage or other liability resulting from use of Gallery 75 Dry Flowable on plant species not recommended on this label.**

Note: Do not apply Gallery 75 Dry Flowable to the following plant species or injury may occur:

Ajuga spp. (bugle)
Brassica spp. (mustard)
Euonymus alatus 'compacta' (dwarf burning bush)
Euphorbia spp. (spurge)
Hydrangea spp. (hydrangea)
Iberi spp. (candytuft)
Juniperus horizontalis 'Prince of Wales' (Prince of Wales Juniper)
Melaleuca quinquenervia (cajeput tree)
Rhododendron carolinianum (Carolina rhododendron)
Rhododendron catawbiense 'Roseum elegans' (Roseum elegans rhododendron)
Sedum spp. (stonecrop)
Yucca recurvifolia (green yucca)

Tank Mixing

Gallery 75 Dry Flowable may be tank mixed with Roundup herbicide or other registered postemergence herbicides to control existing unwanted

vegetation and provide residual preemergence broadleaf weed control in ornamental areas. Gallery 75 Dry Flowable may also be tank mixed with Surflan A.S. herbicide and applied preemergence to provide broad spectrum control of annual grasses and broadleaf weeds in ornamental areas. Applied as directed, Gallery 75 Dry Flowable tank mixes will provide control of susceptible weed species listed on the respective labels. When using Gallery 75 Dry Flowable in tank mix combination with other products, read and follow all applicable use directions, precautions, and limitations on the respective product labels.

Note: Do not apply sprays containing Roundup over the top of ornamental plants. Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying Roundup in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage or stems of desirable plants.

Gallery 75 Dry Flowable may be used on the following established plant species:

(Note: Limitations on recommended treatment methods)

Trees

Scientific Name/Common Name	Recommended Treatment Method: C = Container Grown F = Field Grown
<i>Abies concolor</i>	
White fir	F
<i>Abutilon hybridum</i>	
Albus-flowering maple	F
Luteus-flowering maple	F
Roseus-flowering maple	F
Tangerine-flowering maple	F
Vesuvius red-flowering maple	F
<i>Acer ginnala</i>	
Flame maple	F
<i>Acer rubrum</i>	
Red maple	F
Red sunset maple	F
<i>Acer saccharinum</i>	
Silver maple	C, F
<i>Areacastrum romanzoffianum</i>	
Queen palm	C, F
<i>Betula nigra</i>	
River birch	C, F
<i>Betula papyrifera</i>	
Paper birch	F
<i>Brachychiton populneus</i>	
Bottle tree	C, F
<i>Bucida buceras</i>	
Black olive	F
<i>Ceratonia siliqua</i>	
Carob	F
<i>Cercis canadensis</i>	
Redbud	C, F
<i>Chamaecyparis obtusa</i>	
Filicoides-fernspray cypress	F
Gracilis-slender Hinoki cypress	F
<i>Chamaecyparis pisifera</i>	
Sawara-false cypress	F
Squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	
Palm	F
<i>Chamaedorea costaricana</i>	
Palm	F
<i>Chamaedorea elegans</i>	
Parlor palm	F

<i>Cornus florida</i>	
Cloud nine dogwood	C, F
Flowering dogwood	C, F
<i>Crataegus viridis</i>	
Green hawthorn	F
<i>Cupaniopsis anacardioides</i>	
Carrot wood	F
<i>Cupressus glabra</i>	
Arizona cypress	F
<i>Elaeagnus angustifolia</i>	
Russian olive	C, F
<i>Eucalyptus camaldulensis</i>	
Red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	
Mealy eucalyptus	F
Silver dollar eucalyptus	F
<i>Eucalyptus microtheca</i>	
Coolibah tree	C, F
<i>Eucalyptus sideroxylon</i>	
Red ironbark eucalyptus	F
<i>Ficus benjamina</i>	
Ficus	C, F
Mini ficus	C, F
<i>Fraxinus udhei</i>	
Shamel ash	C, F
<i>Ginkgo biloba</i>	
Maidenhair tree	F
<i>Gleditsia triacanthos</i>	
Honey locust	F
Shademaster honey locust	F
<i>Heteromeles arbutiflora</i>	
Toyon	F
<i>Illicium floridanum</i>	
Florida anise-tree	C, F
<i>Juniperus virginiana</i>	
Eastern red cedar	C, F
<i>Liquidambar styraciflua</i>	
American sweet gum	F
<i>Magnolia grandiflora</i>	
Southern magnolia	C, F
<i>Morus alba</i>	
White mulberry	F
<i>Musa aluminata</i>	
Banana	C, F
<i>Oxydendrum arboreum</i>	
Sourwood	C, F
<i>Picea abies</i>	
Norway spruce	F
Pendula-weeping Norway spruce	F
Repens-spreading Norway spruce	F
<i>Picea glauca conica</i>	
Dwarf Alberta spruce	F
<i>Picea pungens</i>	
Dwarf globe blue spruce	C, F
Glauca-Colorado blue spruce	F
Hoopsii-Hoop's blue spruce	F
Koster-Koster blue spruce	F
<i>Pinus aristata</i>	
Bristlecone pine	F
<i>Pinus canariensis</i>	
Canary Island pine	F
<i>Pinus contorta</i>	
Shore pine, beach pine	F
<i>Pinus eldarica</i>	
Eldarica pine	C, F
<i>Pinus leucodermis</i>	
Bosnian pine	C, F
<i>Pinus mugo</i>	
Pumilio-shrubby swiss mountain pine	C, F

<i>Pinus nigra</i>		<i>Acer palmatum</i>	
Austrian black pine	C, F	Coral bark Japanese maple	C, F
<i>Pinus radiata</i>		Dwarf Japanese maple	C, F
Monterey pine	F	<i>Agave americana</i>	
<i>Pinus strobus</i>		Century plant	F
Eastern white pine	C, F	<i>Astilbe chinensis</i>	
<i>Pinus sylvestris</i>		False spiraea	C, F
Columnar Scotch pine	C, F	<i>Athyrium nipponicum</i>	
Scotch pine	C, F	Japanese painted fern	C, F
<i>Pinus thunbergiana</i>		<i>Baccharis pilularis</i>	
Japanese black pine	F	Coyotebush	F
<i>Platanus occidentalis</i>		<i>Berberis gladwynensis</i>	
American sycamore	F	William Penn barberry	C, F
<i>Platanus racemosa</i>		<i>Berberis mentorensis</i>	
Californian sycamore	F	Mentor barberry	C, F
<i>Podocarpus</i> spp.		<i>Berberis thunbergii</i>	
Podocarpus	F	Atropurea-redleaf Japanese barberry	F
<i>Populus deltoides</i>		Aurea-golden Japanese barberry	F
Cottonwood	F	Crimson pygmy barberry	C, F
<i>Prosopis chilensis</i>		Rose glow barberry	C, F
Chilean mesquite	C, F	<i>Bougainvillea</i> spp.	
<i>Prunus yedoensis</i>		Barbara Karst	F
Yoshino flowering cherry	F	California gold	F
<i>Quercus ilicifolia</i>		Pink pixie	C, F
Bear oak	F	Scarlet O'Hara	F
<i>Quercus palustris</i>		Temple fire	C, F
Pin oak	F	Texas dawn	F
<i>Quercus phellos</i>		<i>Buxus microphylla japonica</i>	
Willow oak	C, F	Japanese boxwood	C, F
<i>Quercus rubra</i>		<i>Buxus microphylla Koreanana</i>	
Red oak	C, F	Korean boxwood	F
<i>Quercus virginiana</i>		<i>Callistemon citrinus</i>	
Live oak	C, F	Lemon bottlebrush	F
<i>Salix babylonica</i>		<i>Callistemon viminalis</i>	
Babylon weeping willow	F	Weeping bottlebrush	C, F
Corkscrew willow	F	<i>Calluna vulgaris</i>	
<i>Sequoiadendron giganteum</i>		Spring torch scotch heather	C, F
Giant sequoia	F	<i>Camellia japonica</i>	
<i>Swietenia mahogani</i>		Camellia	C, F
Mahogany	F	<i>Cassia artemisioides</i>	
<i>Tabebuia caraiba</i>		Feathery cassia	C, F
Yellow tab	F	<i>Ceanothus</i> spp.	
<i>Tsuga canadensis</i>		Wild lilac	F
Eastern hemlock	C, F	<i>Cephalotaxus drupacea</i>	
<i>Ulmus parvifolia</i>		Plum yew	C, F
Chinese elm	F	<i>Cerastium tomentosum</i>	
<i>Washingtonia robusta</i>		Snow-in-summer	C, F
Mexican fan palm	F	<i>Chamaecyparis obtusa</i>	

Ornamental Shrubs

Scientific Name/Common Name	Recommended Treatment Method: C = Container Grown F = Field Grown		
<i>Abelia grandiflora</i>		<i>Chamaecyparis pisifera</i>	
Edward Goucher abelia	C, F	Filifera-thread cypress	F
Glossy abelia	C, F	<i>Chrysalidocarpus lutescens</i>	
<i>Acacia abyssinica</i>		Areca palm	F
Abyssinica acacia	C, F	<i>Clethra alnifolia</i>	
<i>Acacia redolens</i>		Summersweet	C, F
Prostrate acacia	F	<i>Cleyera japonica</i>	
<i>Acacia stenophylla</i>		Japanese cleyera	C, F
Shoestring acacia	C, F	<i>Cornus alba</i>	
<i>Acalypha wilkesiana</i>		Sibirica-Siberian dogwood	F
Copper leaf	C, F	<i>Cornus stolonifera</i>	
<i>Acer ginnala</i>		Baileyi-red-osier dogwood	F
Amur maple	C, F	Flaviramea-yellowtwig dogwood	F
		<i>Cotinus coggygria</i>	
		Royal purple smoke tree	C, F
		<i>Cotinus dammeri</i>	
		Coral beauty smoke tree	C, F
		Eichholz smoke tree	C, F

<i>Cotoneaster adpressus</i>		<i>Ilex cornuta</i>	
Praecox-early cotoneaster	F	Burford holly	C, F
<i>Cotoneaster apiculatus</i>		Dwarf Burford holly	F
Cranberry cotoneaster	C, F	Needle point holly	C, F
<i>Cotoneaster congestus</i>		<i>Ilex crenata</i>	
Pyrenees cotoneaster	F	Compacta-dwarf Japanese holly	C, F
<i>Cotoneaster dammeri</i>		Convexa holly	C, F
Bearberry cotoneaster	C, F	Dwarf Chinese holly	C, F
<i>Cotoneaster himalayan</i>		Green luster holly	C, F
Himalayan cotoneaster	F	Helleri-Heller's Japanese holly	C, F
<i>Cotoneaster horizontalis</i>		Hetzii's Japanese holly	C, F
Rock cotoneaster	C, F	Stokesii Japanese holly	C, F
<i>Cycas revoluta</i>		<i>Ilex glabra</i>	
Sago palm	C, F	Compacta-compact inkberry holly	C, F
<i>Cytisus praecox</i>		Nordica-inkberry holly	C, F
Hollandia-warminster broom	F	<i>Ilex meserveae</i>	
<i>Cytisus scoparius</i>		Blue girl holly	F
Lena-Scotch broom	F	Blue boy holly	F
<i>Daphne odora</i>		Ebony magic holly	F
Fragrant daphne	C, F	<i>Ilex vomitoria</i>	
<i>Deutzia crenata</i>		Nana-dwarf yaupon holly	C, F
Nakiana-dwarf deutzia	F	Pendula-weeping yaupon holly	C, F
<i>Deutzia gracilis</i>		Yaupon holly	C, F
Slender gracilis	F	<i>Illicium annisatum</i>	
<i>Dodonea viscosa</i>		Mystery gardenia	C, F
Hopseed bush	F	<i>Itea ilicifolia</i>	
<i>Elaeagnus pungens</i>		Henry Garnet holly leaf sweetspire	C, F
Fruitland silver berry	C, F	<i>Ixora collinea</i>	
<i>Erica cinerea</i>		Ixora	C, F
Purple bell heather	C, F	<i>Juniperus chinensis</i>	
<i>Erica x darleyensa</i>		Torulosa-hollywood juniper	C, F
Mediterranean pink heather	C, F	Media-old gold juniper	C, F
<i>Erica vagans</i>		Pfitzerana-pfitzer juniper	C, F
Cornish heather	C, F	Pfitzerana glauca-blue juniper	C, F
<i>Euonymus fortunei</i>		Sea green juniper	F
Canadale gold euonymus	F	<i>Juniperus conferta</i>	
Emerald'n gold euonymus	F	Emerald sea shore juniper	C, F
Sunspot euonymus	C, F	Shore juniper	C, F
<i>Euonymus japonica</i>		<i>Juniperus horizontalis</i>	
Silver king euonymus	F	Plumosa-Andorra juniper	C, F
Variegated evergreen euonymus	C, F	Bar Harbor juniper	C, F
<i>Euonymus kiatschovica</i>		Blue rug juniper	C, F
Spreading euonymus	F	Creeping juniper	C, F
<i>Forsythia intermedia</i>		Dwarf Andorra juniper	C, F
Border forsythia	C, F	Huntington blue juniper	C, F
<i>Gardenia jasminoides</i>		Wiltonii-blue carpet juniper	C, F
August beauty gardenia	C, F	<i>Juniperus procumbens</i>	
Gardenia	C, F	Nana-dwarf Japaneses garden juniper	C, F
Radican gardenia	C, F	<i>Juniperus prostrata</i>	
<i>Gaultheria shallon</i>		Prostrata juniper	C, F
Salal/lemon leaf	C, F	<i>Juniperus sabina</i>	
<i>Gelsemium sempervirens</i>		Broadmoor juniper	F
Carolina jessamine	C, F	Foemina-Hicks juniper	C, F
<i>Genista pilosa</i>		Savin juniper	C, F
Woadwaxen	F	Tamariscifolia-Tam juniper	F
<i>Hibiscus rosa-sinensis</i>		<i>Juniperus scopulorum</i>	
Ross Estey-hibiscus	C, F	Emerald green juniper	F
<i>Hibiscus syriacus</i>		<i>Juniperus squamata</i>	
Rose of Sharon-heart	F	Blue juniper	F
Rose of Sharon-red bird	F	Blue star juniper	F
Rose of Sharon-woodbridge	F	Parsonii juniper	C, F
<i>Ilex aquifolium</i>		<i>Kalmia latifolia</i>	
Balkans holly	F	Mountain laurel	C, F
Gold coast holly	F	<i>Lagerstroemia indica</i>	
<i>Ilex aquipernyi</i>		Crape myrtle	C, F
San Jose holly	C, F	<i>Lantana spp.</i>	
<i>Ilex attenuata</i>		Lantana	C, F
Savannah holly	C, F	<i>Leucothoe axillaris</i>	
		Coast leucothoe	C, F

<i>Leucothoe fontainesiana</i>			<i>Potentilla fragiformis</i>	
Drooping leucothoe	C, F		Cinquefoil	F
<i>Ligustrum lucidum</i>			<i>Potentilla fruticosa</i>	
Glossy privet	C, F		Cinquefoil	C, F
<i>Ligustrum ovalifolium</i>			Gold drop potentilla	F
California privet	F		Goldfinger potentilla	C, F
<i>Ligustrum japonicum</i>			Red ace potentilla	C, F
Japanese privet	C, F		Sunset potentilla	C, F
Wax ligustrum	C, F		Tangerine potentilla	C, F
Yellow tip ligustrum	C, F		<i>Potentilla verna</i>	
<i>Ligustrum texanum</i>			Spring cinquefoil	C, F
Howardi privet	F		<i>Prunus glandulosa</i>	
Wax leaf privet	C, F		Dwarf pink flowering almond	C, F
<i>Ligustrum vicaryi</i>			<i>Pyracantha fortuneana</i>	
Vicary golden privet	F		Lalendel Monrovia pyracantha	F
<i>Ligustrum vulgare</i>			Monon pyracantha	F
Lodense privet	C, F		Red elf hybrid pyracantha	F
<i>Livistona chinensis</i>			Rutgers hybrid pyracantha	C, F
Chinese fountain palm	F		Santa Cruz pyracantha	F
<i>Lonicera periclymenum</i>			Victory pyracantha	F
Flowering woodbine	F		<i>Rhaphiolepis indica</i>	
Serotina woodbine	F		Charisma-Monruce raphiolepis	F
<i>Lonicera sempervirens</i>			Enchantress-Moness raphiolepis	F
Trumpet honeysuckle	F		India hawthorn	C, F
<i>Mahonia bealei</i>			Springtime-Monme raphiolepis	F
Leather leaf mahonia	C, F		<i>Rhaphiolepis ovata</i>	
<i>Mahonia repens</i>			Roundleaf raphiolepis	
Creeping mahonia	C, F		Rhododendron spp. (including azalea)	
<i>Myrica cerifera</i>			Azaleas	
Wax myrtle	C, F		<i>Rhododendron calendulacem</i>	
<i>Nandina domestica</i>			Flame azalea	F
Compacta-dwarf heavenly bamboo	C, F		Cannon's double azalea	C, F
Harbour dwarf-heavenly bamboo	C, F		Golden flare azalea	C, F
Heavenly bamboo	C, F		Klondike azalea	C, F
Nana compacta-heavenly bamboo	C, F		<i>Rhododendron indica</i>	
Nana purpurea-heavenly bamboo	C, F		Formosa azalea	C, F
Woods dwarf-heavenly bamboo	C, F		Waucabusa azalea	C, F
<i>Nerium oleander</i>			<i>Rhododendron kaempferi</i>	
Hardy red oleander	F		Blue danube azalea	C, F
Oleander	C, F		<i>Rhododendron kerume</i>	
Ruby lace oleander	F		Coral bells azalea	C, F
<i>Osmanthus fortunei</i>			Hino pink azalea	C, F
Fortunes osmanthus	C, F		Mildred azalea	C, F
<i>Phoenix roeloenii</i>			Snow azalea	C, F
Pigmy date palm	C, F		<i>Rhododendron obtusum</i>	
<i>Photinia fraseri</i>			Hino-crimsom azalea	C, F
Fraser's photinia	C, F		<i>Rhododendron satsumi</i>	
<i>Pieris japonica</i>			Gumpo pink azalea	C, F
Lily-of-the-valley	C, F		Higasa azalea	F
Mountain fire lily-of-the-valley	C, F		Reijn azalea	C, F
Snowdrift lily-of-the-valley	C, F		<i>Rhododendron</i> spp. hybrids	
Temple bells lily-of-the-valley	C, F		Carror azalea	C, F
Valley rose lily-of-the-valley	C, F		Fashion azalea	C, F
Valley valentine lily-of-the-valley	C, F		Gerard Christina azalea	F
<i>Pieris japonica x forestii</i>			Girard Roberta azalea	F
Forest flame lily-of-the-valley	C, F		Helmut vogel azalea	F
<i>Pinus mugo</i>			Hersey red azalea	F
Mugo-mugho pine	C, F		Hot shot azalea	C, F
<i>Pittosporum tobira</i>			Hume azalea	F
Wheeler's dwarf pittosporum	C, F		Inga azalea	F
Green pittosporum	C, F		Irene Koster azalea	C, F
<i>Plumbago ariculata</i>			President Clay azalea	C, F
Blue cape plumbago	F		Tradition azalea	
<i>Plumbago capensis</i>			Rhododendrons	
Plumbago	C, F		<i>Rhododendron campylocarpum</i>	
<i>Podocarpus macrophyllus</i>			Butterfly rhododendron	F
Yewpine	C, F		<i>Rhododendron carolinianum x daurium</i>	
<i>Polystichum polyblepharum</i>			PJM rhododendron	F
Tassel fern	C, F			

<i>Rhododendron catawbiense</i>		<i>Thuja orientalis</i>	
Catawba rhododendron	C, F	Aureus nana-dwarf golden arborvitae	F
Catawba album rhododendron	C, F	Minima glauca-dwarf arborvitae	F
Lord Roberts rhododendron	C, F	<i>Veitchia merrilli</i>	
Rocket rhododendron	C, F	Christmas palm	F
<i>Rhododendron caucasicum x ponticum</i>		<i>Viburnum bodnantense</i>	
Cunninham White rhododendron	F	Pink dawn viburnum	C, F
<i>Rhododendron forrestii x griersonianum</i>		<i>Viburnum davidii</i>	
Elizabeth rhododendron	C, F	David viburnum	F
<i>Rhododendron forrestii repens</i>		<i>Viburnum japonicum</i>	
Gomer Waterer rhododendron	C, F	Viburnum	F
<i>Rhododendron griffithianum</i>		<i>Viburnum lantana</i>	
Jean Marie rhododendron	F	Wayfaring tree	F
<i>Rhododendron impeditum</i>		<i>Viburnum opulus sterile</i>	
Rhododendron	C, F	Common snowball viburnum	F
<i>Rhododendron maximum</i>		<i>Viburnum plicatum tomentosum</i>	
Rhodie max (rosebay)	F	Doublefile viburnum	F
<i>Rhododendron mucronulatum</i>		<i>Viburnum x pragense</i>	
Rhododendron	F	Viburnum	F
<i>Rhododendron ponticum</i>		<i>Viburnum setigerum</i>	
Chionoides rhododendron	F	Tea viburnum	F
Daphnoides rhododendron	C, F	<i>Viburnum tinus compactum</i>	
<i>Rhododendron racemosum</i>		Spring bouquet viburnum	F
Dwarf scarlet wonder rhododendron	C, F	<i>Viburnum trilobum</i>	
Tribly rhododendron	C, F	Cranberry bush	C, F
Unique rhododendron	C, F	<i>Viburnum trilobum compactum</i>	
Vulcan rhododendron	C, F	Dwarf cranberry bush	C, F
<i>Rhododendron sassthigiatum x carolinianum</i>		<i>Weigela florida</i>	
Ramapo rhododendron	F	Bristol ruby weigela	F
<i>Rhododendron hybrid spp.</i>		Java red weigela	F
America rhododendron	C, F	Minuet weigela	F
English Roseum rhododendron	F	<i>Xylosma congestum</i>	
Nova Zembla rhododendron	C, F	Xylosma	F
Scintillation rhododendron	C, F	<i>Yucca filamentosa</i>	
<i>Rhus lancea</i>		Yucca	C, F
Africa sumac	C, F		
<i>Rosa rugosa</i>		Ground Covers	
Ramanas rose	F		
<i>Rosmarinus officinalis</i>			
Rosemary	F		
<i>Skimmia japonica</i>		Scientific Name/Common Name	
Japanese skimmia	C, F	<i>Ammophila breviligulata</i>	
<i>Skimmia revesiana</i>		Beechgrass	C, F
Reeve's skimmia	C, F	<i>Arctotheca calendula</i>	
<i>Spiraea bumalda</i>		Cape weed	F
Anthony Waterer spiraea	C, F	<i>Carex spp.</i>	
<i>Spiraea japonica</i>		Variegated carex	C, F
Dolchia spiraea	C, F	<i>Carpobrotus edulis</i>	
Japanese alpine spiraea	C, F	Largeleaf iceplant	F
Shirobana spiraea	C, F	<i>Cortaderia selloana</i>	
<i>Spiraea vanhouttei</i>		Pampas grass	C, F
Bridal wreath	C, F	<i>Delosperma alba</i>	
<i>Syringa rothomagensis</i>		White iceplant	F
Chinese lilac	C, F	<i>Descampsia caespitosa</i>	
<i>Syringa vulgaris</i>		Descampsia	C, F
Common lilac	F	<i>Drosanthemum floribundum</i>	
<i>Taxus cuspidata</i>		Trailing rosea iceplant	F
Japanese yew	F	<i>Drosantheumum hispidum</i>	
<i>Tecomaria capensis</i>		Iceplant	C, F
Cape honeysuckle	C, F	<i>Festuca ovina glauca</i>	
<i>Ternstroemia gymnanthera</i>		Blue fescue	C, F
<i>Thuja occidentalis</i>		<i>Gazania spp.</i>	
Emerald arborvitae	F	Gazania	C, F
Globosa-globe arborvitae	C, F	<i>Hakonechloa macroaureola</i>	
Little giant-dwarf arborvitae	F	Golden hakonechloa	C, F
Nigra-dark American arborvitae	F	<i>Hedera canariensis</i>	
Pyramidalis-pyramid arborvitae	F	Algerian ivy	F
Rheingold arborvitae	F	<i>Hedera helix</i>	
Techny arborvitae	F	English ivy	C, F

<i>Hemerocallis</i> spp.	
Daylily	C, F
<i>Hosta lancifolia</i>	
Narrow leafed plantain lily	C
<i>Hosta lancifolia</i>	
Albo-marginata hosta	C, F
<i>Hypericum</i> spp.	
St. Johnswort	F
<i>Jasminum nitidum</i>	
Angelwing jasmine	C, F
<i>Lampranthus spectabilis</i>	
Trailing iceplant	F
<i>Liriope gigantea</i>	
White lily turf	F
<i>Liriope muscari</i>	
Big blue lily turf	C, F
Lilac beauty lily turf	C, F
Majestic lily turf	C, F
Monroe white lily turf	C, F
Silvery sunproof lily turf	C, F
Variegated liriope lily turf	C, F
<i>Liriope spicata</i>	
Green/creeping lily turf	C, F
<i>Lonicera japonica</i>	
Japanese honeysuckle	F
<i>Miscanthus sinensis</i>	
Eulalia grass	C, F
<i>Ophiopogon japonicus</i>	
Dwarf Mondo grass	C, F
Mondo grass	C, F
<i>Osteospermum fruticosum</i>	
Trailing African daisy	F
<i>Pachysandra terminalis</i>	
Japanese spurge	C, F
<i>Pennisetum alopecuroides</i>	
Fountain grass	C, F
<i>Phalaris arundinacea picta</i>	
Ribbon grass	C, F
<i>Trachelospermum asiaticum</i>	
Asaian jasmine	C, F
<i>Vinca minor</i>	
Dwarf periwinkle	F

Ornamental Bulbs

Gallery 75 Dry Flowable may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply Gallery 75 Dry Flowable to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. Gallery 75 Dry Flowable may also be applied following bulb emergence but prior to flowering and after flowering. For fall planted bulbs, apply Gallery 75 Dry Flowable again in late winter or early spring to weed-free soil surfaces.

Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.

Do not apply to gladiolus prior to emergence or if corms are less than one inch in diameter.

Do not apply to bulbs while they are flowering.

Non-bearing† Fruit and Nut Trees and Non-bearing† Vineyards

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Common Name

almond	F
apple	F
apricot	F

avocado	F
blackberry	F
blueberry	F
boysenberry	F
cherry, sour	F
cherry, sweet	F
currant	F
dewberry	F
elderberry	F
fig	F
filbert	F
gooseberry	F
grape, American	F
grape, European	F
grapefruit	F
kiwi	F
lemon	F
loganberry	F
macadamia nut	F
nectarine	F
olive F	
orange	F
peach	F
pear F	
pecan	F
pistachio	F
plumF	
prune	F
pomegranate	F
raspberry	F
walnut, black	F
walnut, English	F

†Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

Christmas Tree Plantations

Gallery 75 Dry Flowable Alone

Apply Gallery 75 Dry Flowable as a directed spray to the soil surface or as an overtop spray to established plantings of field grown Christmas tree species listed in this label. Do not apply to seedbeds or seedling transplant beds. Apply only to established plantings. Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation. Follow all instructions provided in the "General Information" section of this label.

Tank Mix Combinations

Tank mix combinations of Gallery 75 Dry Flowable plus other labeled herbicides may be used as directed or overtop sprays in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Gallery 75 Dry Flowable Plus Roundup: Apply tank mix combinations of Gallery 75 Dry Flowable plus Roundup only as directed sprays in Christmas tree plantings. When applied according to use directions, Gallery 75 Dry Flowable plus Roundup herbicide will provide postemergence control of susceptible weed species listed on the label for Roundup and residual preemergence control of susceptible weed species listed on the label for Gallery 75 Dry Flowable. Refer to the label for Roundup for specific use directions, precautions and limitations before use.

Precautions:

Do not apply sprays containing Roundup over the top of Christmas tree plantings.

Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of Christmas trees or other desirable plants or severe damage or death may result.

Non-cropland Areas

Gallery 75 Dry Flowable is recommended as a preemergence herbicide for control of certain broadleaf weeds in non-cropland areas, including industrial sites, utility substations, highway guard rails, sign posts and delineators.

Apply Gallery 75 Dry Flowable anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the "General Information" section prior to using this product on non-cropland.

Gallery 75 Dry Flowable is compatible and can be tank mixed with other herbicides registered for use on non-cropland. Applied as directed, tank mixes containing Gallery 75 Dry Flowable will provide control of susceptible weed species listed on the respective labels. All directions, precautions and limitations on the respective product labels apply to the tank mix use.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

*Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Label Code: D02-081-008

Replaces Label: 113-78-007

EPA-accepted 10/01/93

Revisions:

DowElanco notified the EPA that its legal name will change to Dow AgroSciences LLC effective January 1, 1998. The following label changes are being made via non-notification as a result of this company name change:

- Company name changed from DowElanco to Dow AgroSciences LLC (logo, trademark reference, address line, warranty statement)
- Emergency telephone number updated to include a web site address (PR Notice 97-4)

The name of this product and its EPA registration number remain the same. No other changes are being made to the label at this time.